

L1 – 6.1/6.2 – Intro to Logarithms and Review of Exponentials

MHF4U

Jensen

In this section you will learn about how a logarithmic function is the inverse of an exponential function. You will also learn how to express exponential equations in logarithmic form.

Part 1: Review of Exponential Functions

Equation: $y = a(b)^x$

a = initial amount

b = growth ($b > 1$) or decay ($0 < b < 1$) factor

y = future amount

x = number of times a has increased or decreased

To calculate x , use the equation: $x = \frac{\text{total time}}{\text{time it takes for one growth or decay period}}$

Example 1: An insect colony has a current population of 50 insects. Its population doubles every 3 days.

a) What is the population after 12 days?


b) How long until the population reaches 25 600?

Part 2: Review of Inverse Functions

Inverse of a function:

- The inverse of a function f is denoted as f^{-1}
- The function and its inverse have the property that if $f(a) = b$, then $f^{-1}(b) = a$
- So if $f(5) = 13$, then $f^{-1}(13) = 5$
- More simply put: The inverse of a function has all the same points as the original function, except that the x 's and y 's have been reversed.

The **graph** of $f^{-1}(x)$ is the graph of $f(x)$ reflected in the line $y = x$. This is true for all functions and their inverses.



Example 2: Determine the equation of the inverse of the function $f(x) = 3(x - 5)^2 + 1$

Algebraic Method for finding the inverse:

1. Replace $f(x)$ with " y "
2. Switch the x and y variables
3. Isolate for y
4. replace y with $f^{-1}(x)$

Equation of inverse:

Part 3: Review of Exponent Laws

Name	Rule
Product Rule	$x^a \cdot x^b =$
Quotient Rule	$\frac{x^a}{x^b} =$
Power of a Power Rule	$(x^a)^b =$
Negative Exponent Rule	$x^{-a} =$
Exponent of Zero	$x^0 =$

Part 4: Inverse of an Exponential Function

Example 3:

a) Find the equation of the inverse of $f(x) = 2^x$.


This step uses the 'change of base' formula that we will cover later in the unit.

$$\log_b m = \frac{\log m}{\log b}$$

b) Graph the both $f(x)$ and $f^{-1}(x)$.

$f(x) = 2^x$	
x	y

$f^{-1}(x) = \log_2 x$	
x	y



Note: just swap x and y coordinates to get key points for the inverse of a function. The graph should appear to be a reflection across the line $y = x$.

c) Complete the chart of key properties for both functions

$y = 2^x$	$y = \log_2 x$
x -int:	x -int:
y -int:	y -int:
Domain:	Domain:
Range:	Range:
Asymptote:	Asymptote:

Part 5: What is a Logarithmic Function?

The logarithmic function is the _____ of the exponential function with the same base.

The **logarithmic function** is defined as $y = \log_b x$, or y equals the logarithm of x to the base b .

The function is defined only for _____

In this notation, _____ is the exponent to which the base, _____, must be raised to give the value of _____.

In other words, the solution to a logarithm is always an _____.

The logarithmic function is most useful for solving for unknown _____

_____ are logarithms with a base of 10. It is not necessary to write the base for common logarithms: $\log x$ means the same as $\log_{10} x$

Part 6: Writing Equivalent Exponential and Logarithmic Expressions

Exponential equations can be written in logarithmic form, and vice versa

$$y = b^x \rightarrow$$

$$y = \log_b x \rightarrow$$

Example 4: Rewrite each equation in logarithmic form

a) $16 = 2^4$

b) $m = n^3$

c) $3^{-2} = \frac{1}{9}$

Example 5: Write each logarithmic equation in exponential form

a) $\log_4 64 = 3$

b) $y = \log x$

Note: because there is no base written, this is understood to be the common logarithm of x .

Part 7: Evaluate a Logarithm

Example 6: Evaluate each logarithm without a calculator

Rule: if $x^a = x^b$, then $a = b$

a) $y = \log_3 81$

Rule: $\log_a(a^b) = b$

a) $y = \log_4 64$

Note: either of the rules presented above are appropriate to use for evaluating logarithmic expressions

b) $y = \log\left(\frac{1}{100}\right)$

c) $y = \log_2\left(\frac{1}{8}\right)$